

Short-Crested Breaking Waves and Vorticity

David B. Clark
Woods Hole Oceanographic Institution, MS12
Woods Hole, MA 02543
phone: (508) 289-2935 fax: (508) 457-2194 email: dclark@whoi.edu

Steve Elgar
Woods Hole Oceanographic Institution, MS11
Woods Hole, MA 02543
phone: (508) 289-3614 fax: (508) 457-2194 email: elgar@whoi.edu

Britt Raubenheimer
Woods Hole Oceanographic Institution, MS12
Woods Hole, MA 02543
phone: (508) 289-3427 fax: (508) 457-2194 email: britt@whoi.edu

Award Number: N00014-12-10511
<http://www.whoi.edu/science/AOPE/people/dclark/>

LONG-TERM GOALS

The long-term goal is to determine the contribution of short-crested breaking waves to vorticity (and thus mixing) in nearshore regions and near strong flows from inlets or river mouths.

OBJECTIVES

The objectives of our research in FY13 were:

- Preparation and revision of a manuscript on vorticity generated by short-crested breaking waves
- To investigate the relationship between vorticity generation and the low frequency eddies that drive surfzone tracer dispersion
- Determine the usefulness of high-resolution ultra wide-angle optical images for measuring breaking wave crest lengths

APPROACH

Our approach is to develop new field methods and instrumentation to enable measurements of the processes affecting vorticity, and to assess the importance of those physical processes to numerical models of the nearshore region.

Report Documentation Page			Form Approved OMB No. 0704-0188		
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE 30 SEP 2013		2. REPORT TYPE		3. DATES COVERED 00-00-2013 to 00-00-2013	
4. TITLE AND SUBTITLE Short-Crested Breaking Waves and Vorticity			5a. CONTRACT NUMBER		
			5b. GRANT NUMBER		
			5c. PROGRAM ELEMENT NUMBER		
6. AUTHOR(S)			5d. PROJECT NUMBER		
			5e. TASK NUMBER		
			5f. WORK UNIT NUMBER		
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) Woods Hole Oceanographic Institution,MS12,Woods Hole,MA,02543			8. PERFORMING ORGANIZATION REPORT NUMBER		
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)			10. SPONSOR/MONITOR'S ACRONYM(S)		
			11. SPONSOR/MONITOR'S REPORT NUMBER(S)		
12. DISTRIBUTION/AVAILABILITY STATEMENT Approved for public release; distribution unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT Same as Report (SAR)	18. NUMBER OF PAGES 4	19a. NAME OF RESPONSIBLE PERSON
a. REPORT unclassified	b. ABSTRACT unclassified	c. THIS PAGE unclassified			

WORK COMPLETED

We finished the revision and publication of an article titled “Vorticity generation by short-crested breaking waves.” The article was published in Geophysical Research Letters, and featured on the December 2012 cover (Figure 1). The data from the 2011 VORTEX pilot study was further analyzed for connections between vorticity generated by short-crested breaking waves and the low frequency horizontal eddies thought to mediate tracer dispersion.




Figure 1. The cover of Geophysical Research Letters featuring short crested waves breaking at Duck, NC. This image is a small subsection of a high-resolution ultra-wide angle image considered for remote sensing of breaking wave crests.

The initial analysis was completed on high-resolution ultra-wide angle optical images of the surfzone acquired in July 2012. The images were compared with existing beach imagery (e.g., ARGUS) for the potential to identify individual breaking wave faces in the surfzone and to measure the lengths of breaking crests (thought to control wave generated vertical vorticity).

Low frequency eddies are thought to be the primary process that disperses tracers in the surfzone and nearshore, but the mechanisms that generate these eddies are not understood. We found that the strength of low frequency eddies was highly correlated with the magnitude of vorticity generated by short crested breaking waves, which varied over a tidal cycle (Figure 2). The high correlation suggests that short-crested breaking waves are a significant control on nearshore tracer dispersion, and the effects of short-crested breaking should be included in wave averaged models when accurate eddy diffusivity or tracer dispersion estimates are required.

3

IMPACT/APPLICATIONS

This project made the first measurements of surfzone vorticity about a vertical axis (i.e., horizontal eddies), and the first measurements of vertical vorticity generated by short crested breaking waves. Short-crested breaking is found to be a significant source of surfzone vorticity, and high correlations with low frequency eddies suggest short-crested breaking is a controlling factor on surfzone tracer dispersion. Our study indicates that the effects of short crested breaking waves should be included in numerical models of the surfzone.

High-resolution ultra wide-angle images of the surfzone are a promising tool for measuring breaking crest-length and other parameters that could determine the influence of short-crested breaking on surfzone flows. The image processing required for this analysis is complex and beyond the scope of this project, however we plan to continue exploring image-processing solutions in the future. An image processing planning letter was submitted in April 2013 to the ONR Littoral Geosciences & Optics Program for CORE funding, with extensions to identifying active wave breaking with radar.

RELATED PROJECTS

An ongoing surfzone vorticity project funded by NSF complements the pilot project analysis funded by this ONR project. A month-long time series of surfzone vorticity will be used to explore the variation in wave-generated vorticity over a range of wave conditions.

PUBLICATIONS

Clark, D.B., S. Elgar, and B. Raubenheimer (2012), (COVER) Vorticity generation by short-crested breaking waves, *Geophysical Research Letters*, 39, L24604, doi:10.1029/2012GL054034, [published, refereed]

This article is available at:

http://www.whoi.edu/science/AOPE/people/dclark/pubs/Clark_etal_GRL_2012.pdf